

| | |
|-------------|---|
| Title | Some Doubly Infinite and Mixed Infinite Sums derived from The N-Fractional Calculus of A Logarithmic Function : with Some Examinations (Coefficient Inequalities in Univalent Function Theory and Related Topics) |
| Author(s) | Nishimoto, Katsuyuki; Romero, Susana S. de; Prieto, Ana I. |
| Citation | 数理解析研究所講究録 (2005), 1414: 15-25 |
| Issue Date | 2005-02 |
| URL | http://hdl.handle.net/2433/26227 |
| Right | |
| Type | Departmental Bulletin Paper |
| Textversion | publisher |

Some Doubly Infinite and Mixed Infinite Sums derived from The N- Fractional Calculus of A Logarithmic Function (with Some Examinations)

Katsuyuki Nishimoto , Susana S. de Romero
and Ana I Prieto

Abstract

In this article theorems for some doubly infinite and mixed infinite sums derived from the N-fractional calculus of a logarithmic function are reported.

Moreover some numerical examinations for the theorems are reported too.

§ 0. Introduction (Definition of Fractional Calculus)

(I) Definition. (by K. Nishimoto) ([1] Vol. 1)

Let $D = \{D_-, D_+\}$, $C = \{C_-, C_+\}$,

C_- be a curve along the cut joining two points z and $-\infty + i\text{Im}(z)$,

C_+ be a curve along the cut joining two points z and $\infty + i\text{Im}(z)$,

D_- be a domain surrounded by C_- , D_+ be a domain surrounded by C_+ .

(Here D contains the points over the curve C).

Moreover, let $f = f(z)$ be a regular function in $D(z \in D)$,

$$f_\nu(z) = (f)_\nu = {}_C(f)_\nu = \frac{\Gamma(\nu+1)}{2\pi i} \int_C \frac{f(\xi)}{(\xi-z)^{\nu+1}} d\xi \quad (\nu \notin \mathbb{Z}), \quad (1)$$

$$(f)_{-m} = \lim_{\nu \rightarrow -m} (f)_\nu \quad (m \in \mathbb{Z}^+), \quad (2)$$

where $-\pi \leq \arg(\xi-z) \leq \pi$ for C_- , $0 \leq \arg(\xi-z) \leq 2\pi$ for C_+ ,

$\xi \neq z$, $z \in C$, $\nu \in \mathbb{R}$, Γ ; Gamma function,

then $(f)_\nu$ is the fractional differintegration of arbitrary order ν (derivatives of order ν for $\nu > 0$, and integrals of order $-\nu$ for $\nu < 0$), with respect to z , of the function f , if $|(f)_\nu| < \infty$.

(II) On the fractional calculus operator N^ν [3]

Theorem A. Let fractional calculus operator (Nishimoto's Operator) N^ν be

$$N^\nu = \left(\frac{\Gamma(\nu+1)}{2\pi i} \int_C \frac{d\xi}{(\xi-z)^{\nu+1}} \right) \quad (\nu \notin \mathbb{Z}), \quad [\text{Refer to (1)}] \quad (3)$$

with

$$N^{-m} = \lim_{\nu \rightarrow -m} N^\nu \quad (m \in \mathbb{Z}^+), \quad (4)$$

and define the binary operation \circ as

$$N^\beta \circ N^\alpha f = N^\beta N^\alpha f = N^\beta(N^\alpha f) \quad (\alpha, \beta \in \mathbb{R}), \quad (5)$$

then the set

$$\{N^\nu\} = \{N^\nu \mid \nu \in \mathbb{R}\} \quad (6)$$

is an Abelian product group (having continuous index ν) which has the inverse transform operator $(N^\nu)^{-1} = N^{-\nu}$ to the fractional calculus operator N^ν , for the function f such that $f \in F = \{f; 0 \neq |f_\nu| < \infty, \nu \in \mathbb{R}\}$, where $f = f(z)$ and $z \in C$. (vis. $-\infty < \nu < \infty$).

(For our convenience, we call $N^\beta \circ N^\alpha$ as product of N^β and N^α .)

Theorem B. " F.O.G. $\{N^\nu\}$ " is an " Action product group which has continuous index ν " for the set of F . (F.O.G. ; Fractional calculus operator group)

Theorem C. Let

$$S := \{\pm N^\nu\} \cup \{0\} = \{N^\nu\} \cup \{-N^\nu\} \cup \{0\} \quad (\nu \in \mathbb{R}). \quad (7)$$

Then the set S is a commutative ring for the function $f \in F$, when the identity

$$N^\alpha + N^\beta = N^\gamma \quad (N^\alpha, N^\beta, N^\gamma \in S) \quad (8)$$

holds. [5]

(III) **Lemma.** We have [1]

$$(i) \quad ((z-c)^\beta)_\alpha = e^{-i\pi\alpha} \frac{\Gamma(\alpha-\beta)}{\Gamma(-\beta)} (z-c)^{\beta-\alpha} \quad \left(\left| \frac{\Gamma(\alpha-\beta)}{\Gamma(-\beta)} \right| < \infty \right),$$

$$(ii) \quad (\log(z-c))_\alpha = -e^{-i\pi\alpha} \Gamma(\alpha) (z-c)^{-\alpha} \quad (|\Gamma(\alpha)| < \infty),$$

$$(iii) \quad ((z-c)^{-\alpha})_{-\alpha} = -e^{i\pi\alpha} \frac{1}{\Gamma(\alpha)} \log(z-c) \quad (|\Gamma(\alpha)| < \infty),$$

where $z-c \neq 0$ in (i), and $z-c \neq 0, 1$ in (ii) and (iii). (Γ ; Gamma function),

$$(iv) \quad (u \cdot v)_\alpha = \sum_{k=0}^{\infty} \frac{\Gamma(\alpha+1)}{k! \Gamma(\alpha+1-k)} u_{\alpha-k} v_k \quad \left(\begin{array}{l} u = u(z), \\ v = v(z) \end{array} \right).$$

§ 1. Doubly Infinite Sum and Mixed One

In the following $\alpha, \beta \in \mathbb{R}$.

Theorem 1. *Let*

$$M(\alpha, \beta; k, m) := \frac{\Gamma(\alpha + k)\Gamma(k + m)\Gamma(\beta + 1)\Gamma(\beta - \alpha - m)}{k! \cdot m! \Gamma(\alpha)\Gamma(k)\Gamma(\beta + 1 - m)\Gamma(-\alpha)}. \quad (1)$$

(i) When $\beta \notin \mathbb{Z}_0^+$, we have the following doubly infinite sums ;

$$\sum_{k=0}^{\infty} \sum_{m=0}^{\infty} M(\alpha, \beta; k, m) \left(\frac{z-c}{z}\right)^m \left(\frac{c}{z}\right)^k \subseteq \frac{\Gamma(\beta - \alpha)}{\Gamma(-\alpha)} \left(\frac{z}{z-c}\right)^{\alpha-\beta}, \quad (2)$$

where

$$z - c \neq 0, 1, \quad z \neq 0, 1, \quad |(z-c)/z| < 1, \quad |c/z| < 1,$$

and

$$|\Gamma(\alpha)|, \quad \left| \frac{\Gamma(\beta - \alpha - m)}{\Gamma(-\alpha)} \right| < \infty.$$

The identity (notation =) holds for $(\alpha - \beta) \in \mathbb{Z}$.

(ii) When $s \in \mathbb{Z}^+$, we have the following mixed infinite sums ;

$$\sum_{k=0}^{\infty} \sum_{m=0}^s M(\alpha, s; k, m) \left(\frac{z-c}{z}\right)^m \left(\frac{c}{z}\right)^k \subseteq \frac{\Gamma(s - \alpha)}{\Gamma(-\alpha)} \left(\frac{z}{z-c}\right)^{\alpha-s}, \quad (3)$$

where

$$z - c \neq 0, 1, \quad z \neq 0, 1, \quad |c/z| < 1, \quad |(z-c)/z| < \infty,$$

and

$$|\Gamma(\alpha)| < \infty.$$

The identity (notation =) holds for $(\alpha - s) \in \mathbb{Z}$.

Proof of (i). We have

$$\log \frac{z-c}{z} = \log \left(1 - \frac{c}{z}\right) = - \sum_{k=1}^{\infty} \frac{c^k}{k} z^{-k} \quad \left(\left| \frac{-c}{z} \right| < 1 \right). \quad (4)$$

Operate N-fractional calculus operator N^α to the both sides of (4), we obtain

$$z^{-\alpha} - (z-c)^{-\alpha} = - \sum_{k=1}^{\infty} \frac{c^k}{k!} \cdot \frac{\Gamma(\alpha + k)}{\Gamma(\alpha)} z^{-k-\alpha} \quad (|\Gamma(\alpha)| < \infty), \quad (5)$$

since

$$N^\alpha \left(\log \frac{z-c}{z} \right) = (\log(z-c) - \log z)_\alpha \quad (z-c \neq 0, 1, z \neq 0, 1) \quad (6)$$

$$= e^{-i\pi\alpha} \Gamma(\alpha) \{z^{-\alpha} - (z-c)^{-\alpha}\} \quad (|\Gamma(\alpha)| < \infty) \quad (7)$$

and

$$N^\alpha (z^{-k}) = (z^{-k})_\alpha = e^{-i\pi\alpha} \frac{\Gamma(k+\alpha)}{\Gamma(k)} z^{-k-\alpha} \quad \left(\left| \frac{\Gamma(k+\alpha)}{\Gamma(k)} \right| < \infty \right) \quad (8)$$

by Lemmas (ii) and (i).

Therefore, we have

$$(z-c)^\alpha - z^\alpha = - \sum_{k=1}^{\infty} \frac{c^k}{k!} \cdot \frac{\Gamma(\alpha+k)}{\Gamma(\alpha)} (z-c)^\alpha z^{-k}, \quad (9)$$

hence

$$z^\alpha = \sum_{k=0}^{\infty} \frac{c^k}{k!} \cdot \frac{\Gamma(\alpha+k)}{\Gamma(\alpha)} (z-c)^\alpha z^{-k}, \quad (10)$$

from (5).

Next operate N-fractional calculus operator N^β to the both sides of (10), we obtain

$$(z^\alpha)_\beta = \sum_{k=0}^{\infty} \frac{c^k}{k!} \cdot \frac{\Gamma(\alpha+k)}{\Gamma(\alpha)} ((z-c)^\alpha z^{-k})_\beta, \quad (11)$$

$$(z^\alpha)_\beta = e^{-i\pi\beta} \frac{\Gamma(\beta-\alpha)}{\Gamma(-\alpha)} z^{\alpha-\beta} \quad \left(\left| \frac{\Gamma(\beta-\alpha)}{\Gamma(-\alpha)} \right| < \infty \right), \quad (12)$$

$$((z-c)^\alpha z^{-k})_\beta = \sum_{m=0}^{\infty} \frac{\Gamma(\beta+1)}{m! \Gamma(\beta+1-m)} ((z-c)^\alpha)_{\beta-m} (z^{-k})_m. \quad (13)$$

Now we have

$$((z-c)^\alpha)_{\beta-m} = e^{-i\pi(\beta-m)} \frac{\Gamma(\beta-\alpha-m)}{\Gamma(-\alpha)} (z-c)^{\alpha-\beta+m} \quad \left(\left| \frac{\Gamma(\beta-\alpha-m)}{\Gamma(-\alpha)} \right| < \infty \right), \quad (14)$$

and

$$(z^{-k})_m = e^{-i\pi m} \frac{\Gamma(k+m)}{\Gamma(k)} z^{-k-m}, \quad (15)$$

by Lemma (iv) and (i), respectively.

Therefore, we obtain

$$\begin{aligned} & \frac{\Gamma(\beta - \alpha)}{\Gamma(-\alpha)} z^{\alpha - \beta} \\ &= \sum_{k=0}^{\infty} \frac{\Gamma(\alpha + k)}{k! \Gamma(\alpha)} \sum_{m=0}^{\infty} \frac{\Gamma(k + m) \Gamma(\beta + 1) \Gamma(\beta - \alpha - m)}{m! \Gamma(k) \Gamma(\beta + 1 - m) \Gamma(-\alpha)} \left(\frac{z - c}{z} \right)^m \left(\frac{c}{z} \right)^k (z - c)^{\alpha - \beta} \quad (16) \end{aligned}$$

from (10) ~ (15).

Therefore, we have

$$\sum_{k=0}^{\infty} \sum_{m=0}^{\infty} M(\alpha, \beta; k, m) \left(\frac{z - c}{z} \right)^m \left(\frac{c}{z} \right)^k = \frac{\Gamma(\beta - \alpha)}{\Gamma(-\alpha)} \left(\frac{z}{z - c} \right)^{\alpha - \beta}, \quad (17)$$

from (16).

However the LHS (left hand side) of (17) is always one valued function, on the contrary the RHS (right hand side) of (17) is many valued function for $(\alpha - \beta) \notin \mathbb{Z}$ and one valued one for $(\alpha - \beta) \in \mathbb{Z}$.

Hence we must calculate as

$$\left(\frac{z}{z - c} \right)^{\alpha - \beta} = \left(e^{i2n\pi} \frac{z}{z - c} \right)^{\alpha - \beta} \quad \left(\begin{array}{l} n \in \mathbb{Z} \\ (\alpha - \beta) \notin \mathbb{Z} \end{array} \right) \quad (18)$$

because we are now being in the field of complex analysis.

Moreover, when $(\alpha - \beta) \in \mathbb{Z}$ both of the LHS and RHS of (17) are one valued functions respectively. In this case we have (17) strictly.

Therefore, we obtain (2) from (17), considering (18) finally.

Proof of (ii). Set $\beta = s \in \mathbb{Z}^+$ in (2), we have then (3) clearly, under the conditions.

§ 2. Some Numerical Examinations for Theorem 1

[I] Examination of Theorem 1. (2)

Set

$$c = 1, \quad z = 10, \quad \alpha = 1/4 \quad \text{and} \quad \beta = 1/2$$

in Theorem 1. (2), we obtain

$$\begin{aligned} \sum_{k=0}^{\infty} \sum_{m=0}^{\infty} M(1/4, 1/2; k, m) \left(\frac{1}{10}\right)^k \left(\frac{9}{10}\right)^m \\ \subseteq -\frac{1}{4} \cdot \frac{\Gamma(1/4)}{\Gamma(3/4)} \left(e^{i2n\pi} \frac{9}{10}\right)^{1/4} \quad (n \in \mathbf{Z}) \quad (1) \\ = \begin{cases} -0.72044\dots & (\text{for } n=0) \quad (2) \\ -i0.72044\dots & (\text{for } n=1) \quad (3) \\ 0.72044\dots & (\text{for } n=2) \quad (4) \\ i0.72044\dots & (\text{for } n=3) \quad (5) \end{cases} \end{aligned}$$

When $\alpha, \beta, c, z \in \mathbf{R}$, the left hand side (LHS) of (3) is real, then we must choose (2) and (4) from the set $\{ (2), (3), (4), (5) \}$.

Now we have

$$\begin{aligned} M(1/4, 1/2; 0, 0) \left(\frac{1}{10}\right)^0 \left(\frac{9}{10}\right)^0 & \quad \left(\begin{array}{l} \text{first term of} \\ \text{the LHS of (1)} \end{array} \right) \\ = \frac{\Gamma(1/4)}{\Gamma(-1/4)} = -\frac{1}{4} \cdot \frac{\Gamma(1/4)}{\Gamma(3/4)} < 0. \quad (6) \end{aligned}$$

Then choosing (2) from the set $\{ (2), (4) \}$, since the sign of the double infinite sum of LHS of (1) is decided by the sign of its first term (with $k = m = 0$), when

$$\left| M_{k,m} \left(\frac{1}{10}\right)^k \left(\frac{9}{10}\right)^m \right| > \left| M_{k+1,m+1} \left(\frac{1}{10}\right)^{k+1} \left(\frac{9}{10}\right)^{m+1} \right|, \quad M_{k,m} = M(\alpha, \beta; k, m),$$

we have then

$$\sum_{k=0}^{\infty} \sum_{m=0}^{\infty} M(1/4, 1/2; k, m) \left(\frac{1}{10}\right)^k \left(\frac{9}{10}\right)^m = -0.72044\dots \quad (7)$$

from (2), considering (6).

Indeed we have

$$\text{LHS of (7)} = \sum_{k=0}^{\infty} \frac{\Gamma(\frac{1}{4}+k)}{k! \Gamma(\frac{1}{4})} \left(\frac{1}{10}\right)^k \sum_{m=0}^{\infty} \frac{\Gamma(k+m) \frac{1}{2} \Gamma(\frac{1}{2}) \Gamma(\frac{1}{4}-m)}{m! \Gamma(k) \Gamma(\frac{1}{2}+1-m) \Gamma(-\frac{1}{4})} \left(\frac{9}{10}\right)^m \quad (8)$$

$$\begin{aligned} &= \frac{\Gamma(\frac{1}{4})}{\Gamma(-\frac{1}{4})} \sum_{k=0}^{\infty} \frac{\Gamma(\frac{1}{4}+k)}{\Gamma(\frac{1}{4})} \left(\frac{1}{10}\right)^k + \frac{\Gamma(-\frac{3}{4})}{2! \Gamma(-\frac{1}{4})} \left(\frac{9}{10}\right) \sum_{k=0}^{\infty} \frac{\Gamma(\frac{1}{4}+k) \cdot k}{k! \Gamma(\frac{1}{4})} \left(\frac{1}{10}\right)^k \\ &\quad - \frac{1}{2! 2^2} \cdot \frac{\Gamma(-\frac{7}{4})}{\Gamma(-\frac{1}{4})} \left(\frac{9}{10}\right)^2 \sum_{k=0}^{\infty} \frac{\Gamma(\frac{1}{4}+k) \cdot k(k+1)}{k! \Gamma(\frac{1}{4})} \left(\frac{1}{10}\right)^k \\ &\quad + \frac{3}{3! 2^3} \cdot \frac{\Gamma(-\frac{11}{4})}{\Gamma(-\frac{1}{4})} \left(\frac{9}{10}\right)^3 \sum_{k=0}^{\infty} \frac{\Gamma(\frac{1}{4}+k) \cdot k(k+1)(k+2)}{k! \Gamma(\frac{1}{4})} \left(\frac{1}{10}\right)^k \\ &\quad - \frac{3 \cdot 5}{4! 2^3} \cdot \frac{\Gamma(-\frac{15}{4})}{\Gamma(-\frac{1}{4})} \left(\frac{9}{10}\right)^4 \sum_{k=0}^{\infty} \frac{\Gamma(\frac{1}{4}+k) \cdot k(k+1)(k+2)(k+3)}{k! \Gamma(\frac{1}{4})} \left(\frac{1}{10}\right)^k \end{aligned}$$

+

(9)

$$\begin{aligned} &= \frac{1}{(-4)} \cdot \frac{\Gamma(\frac{1}{4})}{\Gamma(\frac{3}{4})} \left\{ 1 + \frac{1}{40} + \frac{5}{2! 40^2} + \frac{5 \cdot 9}{3! 40^3} + \frac{5 \cdot 9 \cdot 13}{4! 40^4} + \frac{5 \cdot 9 \cdot 13 \cdot 17}{5! 40^5} + \dots \right\} \\ &\quad + \frac{1}{2 \cdot 3} \cdot \frac{\Gamma(\frac{1}{4})}{\Gamma(\frac{3}{4})} \left(\frac{9}{10}\right) \left\{ 0 + \frac{1}{40} + \frac{5}{40^2} + \frac{5 \cdot 9}{2! 40^3} + \frac{5 \cdot 9 \cdot 13}{3! 40^4} + \frac{5 \cdot 9 \cdot 13 \cdot 17}{4! 40^5} + \dots \right\} \\ &\quad + \frac{1}{2! 3 \cdot 7} \cdot \frac{\Gamma(\frac{1}{4})}{\Gamma(\frac{3}{4})} \left(\frac{9}{10}\right)^2 \left\{ 0 + \frac{2}{40} + \frac{3 \cdot 5}{40^2} \right. \\ &\quad \left. + \frac{4(5 \cdot 9)}{2! 40^3} + \frac{5(5 \cdot 9 \cdot 13)}{3! 40^4} + \frac{6(5 \cdot 9 \cdot 13 \cdot 17)}{4! 40^5} + \dots \right\} \\ &\quad + \frac{1}{3 \cdot 7 \cdot 11} \cdot \frac{\Gamma(\frac{1}{4})}{\Gamma(\frac{3}{4})} \left(\frac{9}{10}\right)^3 \left\{ 0 + \frac{2 \cdot 3}{40} + \frac{3 \cdot 4 \cdot (5)}{40^2} \right. \\ &\quad \left. + \frac{4 \cdot 5(5 \cdot 9)}{2! 40^3} + \frac{5 \cdot 6(5 \cdot 9 \cdot 13)}{3! 40^4} + \frac{6 \cdot 7(5 \cdot 9 \cdot 13 \cdot 17)}{4! 40^5} + \dots \right\} \\ &\quad + \frac{5}{2! 3 \cdot 7 \cdot 11 \cdot 15} \cdot \frac{\Gamma(\frac{1}{4})}{\Gamma(\frac{3}{4})} \left(\frac{9}{10}\right)^4 \left\{ 0 + \frac{2 \cdot 3 \cdot 4}{40} + \frac{3 \cdot 4 \cdot 5(5)}{40^2} \right. \\ &\quad \left. + \frac{4 \cdot 5 \cdot 6(5 \cdot 9)}{2! 40^3} + \frac{5 \cdot 6 \cdot 7(5 \cdot 9 \cdot 13)}{3! 40^4} + \frac{6 \cdot 7 \cdot 8(5 \cdot 9 \cdot 13 \cdot 17)}{4! 40^5} + \dots \right\} \end{aligned}$$

$$+ \dots\dots\dots (10)$$

$$= - (0.75941\dots) + (0.01265\dots) + (0.00205\dots) \\ + (0.00182\dots) + (0.00117\dots) + \dots\dots (11)$$

$$= - 0.74172\dots . (12)$$

[II] Examination of Theorem 1. (3) for $(\alpha - s) \notin \mathbf{Z}$

Set

$$c = 1, \quad z = 3, \quad \alpha = 1/2 \quad \text{and} \quad s = 1$$

in Theorem 1. (3), we obtain

$$\sum_{k=0}^{\infty} \sum_{m=0}^1 M(1/2, 1; k, m) \left(\frac{1}{3}\right)^k \left(\frac{2}{3}\right)^m \\ \subseteq -\frac{1}{2} \left(e^{i 2n\pi} \frac{2}{3} \right)^{1/2} \quad (n \in \mathbf{Z}) \quad (13)$$

$$= \begin{cases} -0.408240\dots & (\text{for } n = 0) \end{cases} \quad (14)$$

$$= \begin{cases} 0.408240\dots & (\text{for } n = 1) \end{cases} \quad (15).$$

Now we have

$$M(1/2, 1; 0, 0) \left(\frac{1}{3}\right)^0 \left(\frac{2}{3}\right)^0 \quad \left(\begin{array}{l} \text{first term of} \\ \text{the LHS of (13)} \end{array} \right) \\ = \frac{\Gamma(1/2)}{\Gamma(-1/2)} = -\frac{1}{2} < 0. \quad (16)$$

Then choosing (14) from the set $\{ (14), (15) \}$, since the sign of the infinite mixed sum of LHS of (13) is decided by the sign of its first term (with $k = m = 0$), when

$$\left| M_{k,m} \left(\frac{1}{3}\right)^k \left(\frac{2}{3}\right)^m \right| > \left| M_{k+1,m+1} \left(\frac{1}{3}\right)^{k+1} \left(\frac{2}{3}\right)^{m+1} \right|, \quad M_{k,m} = M(1/2, 1; k, m),$$

we have then

$$\sum_{k=0}^{\infty} \sum_{m=0}^1 M(1/2, 1; k, m) \left(\frac{1}{3}\right)^k \left(\frac{2}{3}\right)^m = - 0.408240\dots \quad (17)$$

from (14), considering (16).

Indeed we have

$$\text{LHS of (17)} = \sum_{k=0}^{\infty} \frac{\Gamma(\frac{1}{2}+k)}{k! \Gamma(\frac{1}{2})} \left(\frac{1}{3}\right)^k \left(-\frac{1}{2} + k \cdot \frac{2}{3}\right) \quad (18)$$

$$= -\frac{1}{2} + \frac{1}{2 \cdot 3} \cdot \frac{1}{6} + \frac{1}{2!} \cdot \frac{1}{2^2 \cdot 3} \cdot \frac{5}{6} + \frac{1}{3!} \cdot \frac{1}{2^3 \cdot 3^2} \cdot \frac{9}{6} \\ + \frac{1}{4!} \cdot \frac{5 \cdot 7}{2^4 \cdot 3^3} \cdot \frac{13}{6} + \frac{1}{5!} \cdot \frac{5 \cdot 7 \cdot 9}{2^5 \cdot 3^4} \cdot \frac{17}{6} + \dots \quad (19)$$

$$= -0.5 + (0.02777\dots) + (0.034722\dots) + (0.017361\dots) \\ + (0.007314\dots) + (0.002869\dots) + (0.0001083\dots) + \dots \quad (20)$$

$$= -0.40887\dots \quad (21)$$

[III] Examination of Theorem 1. (3) for $(\alpha - s) \in \mathbb{Z}$

Set

$$c = 1, \quad z = 3, \quad \alpha = 2 \quad \text{and} \quad s = 1$$

in Theorem 1. (3), we obtain

$$\sum_{k=0}^{\infty} \sum_{m=0}^1 M(2, 1; k, m) \left(\frac{1}{3}\right)^k \left(\frac{2}{3}\right)^m = \frac{\Gamma(-1)}{\Gamma(-2)} \left(\frac{3}{2}\right) = -3, \quad (22)$$

without the ad hoc shown in [I] and [II] in which the RHS of (1) and (13) are many valued ones.

That is, in this case both sides of § 1. (3) are one valued functions respectively, then we have the notation = in § 1. (3).

Indeed we have

$$\text{LHS of (22)} = \sum_{k=0}^{\infty} \frac{\Gamma(2+k)}{k! \Gamma(k)} \left(\frac{1}{3}\right)^k \sum_{m=0}^1 \frac{\Gamma(k+m) \Gamma(-1-m)}{m! \Gamma(2-m) \Gamma(-2)} \left(\frac{2}{3}\right)^m \quad (23)$$

$$= -2 \sum_{k=0}^{\infty} (k+1) \left(\frac{1}{3}\right)^k + \frac{2}{3} \sum_{k=0}^{\infty} k(k+1) \left(\frac{1}{3}\right)^k \quad (24)$$

$$\begin{aligned}
&= -2 \left(1 + \frac{2}{3} + \frac{3}{3^2} + \frac{4}{3^3} + \frac{5}{3^4} + \dots \right) \\
&\quad + \frac{2}{3} \left(0 + \frac{2}{3} + \frac{2 \cdot 3}{3^2} + \frac{3 \cdot 4}{3^3} + \frac{4 \cdot 5}{3^4} + \frac{5 \cdot 6}{3^5} + \dots \right) \quad (25)
\end{aligned}$$

$$= -2(2.2478\dots) + \frac{2}{3}(2.2313\dots) \quad (26)$$

$$= -3.00807\dots \quad (27)$$

§ 3. Commentary

1. Notice that the LHS of § 1. (2) is always one valued function, on the contrary its RHS is many valued function for $(\alpha - \beta) \notin \mathbb{Z}$ and one valued one for $(\alpha - \beta) \in \mathbb{Z}$.

And notice that when both of the LHS and the RHS of § 1. (2) are one valued functions respectively, namely in the case of $(\alpha - \beta) \in \mathbb{Z}$, we have the identity (notation =) in § 1. (2) always.

2. " Fractional calculus " is essentially a problem in the field of complex analysis. We should not forget that we are now being in the field of fractional calculus, that is, in that of complex analysis.

References

- [1] K. Nishimoto ; Fractional Calculus, Vol. 1 (1984), Vol. 2 (1987), Vol. 3 (1989), Vol. 4 (1991), Vol. 5, (1996), Descartes Press, Koriyama, Japan.
- [2] K. Nishimoto ; An Essence of Nishimoto's Fractional Calculus (Calculus of the 21st Century); Integrals and Differentiations of Arbitrary Order (1991), Descartes Press, Koriyama, Japan.
- [3] K. Nishimoto ; On Nishimoto's fractional calculus operator N^ν (On an action group), J. Frac. Calc. Vol. 4, Nov. (1993), 1 - 11.
- [4] K. Nishimoto ; Unification of the integrals and derivatives (A serendipity in fractional calculus), J. Frac. Calc. Vol. 6, Nov. (1994), 1 - 14.
- [5] K. Nishimoto ; Ring and Field Produced from The Set of N- Fractional Calculus Operator, J. Frac Calc. Vol. 24, Nov. (2003), 29 - 36.
- [6] K. Nishimoto, Ding- Kuo Chyan, Shy- Der Lin and Shih- Tong Tu ; On some infinite sums derived by N- fractional calculus, J. Frac. Calc. Vol.20 (2001), 91 - 97.
- [7] Pin Yu Wang, Tsu- Chen Wu and Shih- Tong Tu; Some Infinite Sums via N- fractional calculus, J. Frac. Calc. Vol.21, May (2002), 71 - 77.

- [8] Shy- Der Lin, Shih- Tong Tu, Tsai- Ming Hsieh and H.M. Srivastava ; Some Finite and Infinite Sums Associated with the Digamma and Related Functions, J. Frac.Calc.Vol.22, Nov. (2002), 103- 114.
- [9] K. Nishimoto ; N- Fractional Calculus of the Power and Logarithmic Functions and Some Identities (Continue), J. Frac. Calc. Vol.22, Nov. (2002), 59 - 65.
- [10] K. Nishimoto, Susana S. de Romero and Ana I. Prieto ; Some Doubly and Triply Infinite Sums derived from The N- Fractional Calculus of A Logarithmic Function. J. Frac Calc.Vol. 24, Nov. (2003), 59 - 66.
- [11] K. Nishimoto and Susana S. de romero ; Some Multiple Infinite Sums derived from The N- Fractional Calculus of some Power Functions, J. Frac. Calc.Vol.24, Nov. (2003), 67 - 76.
- [12] K. Nishimoto ; Examinations for Some Doubly Infinite, Finite and Mixed Sums, J. Frac. Calc. Vol. 25, May (2004), 25 - 32.
- [13] Shy- Der Lin and H. M. Srivastava ; Fractional Calculus and Its Applications Involving Bilateral Expansions and Multiple Infinite Sums, J. Frac. Calc. Vol. 25, May (2004), 47 - 58.
- [14] K. Nishimoto, Susana S. de Romero and Ana I. Prieto ; Examinations for Some Doubly Infinite Sums derived by Means of N-Fractional Calculus, J. Frac. Calc.Vol.26, Nov. (2004), 1 - 8.
- [15] K. Nishimoto ; Some Multiply Infinite, Mixed and Finite Sums derived from The N- Fractional Calculus of Some Power Functions, J. Frac. Calc.Vol.26, Nov. (2004), 9 - 23.
- [16] K. Nishimoto ; Examinations for Some Doubly Infinite Sums derived from The N- Fractional Calculus of A Logarithmic Function, J. Frac.Calc. Vol. 26, Nov. (2004), 25 - 34.
- [17] K. Nishimoto ; On Some $(q + 1)$ Multiply Infinite Sums $(q \in \mathbb{Z}^+)$ derived from the N- Fractional Calculus of Some Power Functions (Part I), J. Frac.Calc.Vol. 26, Nov. (2004), 53 - 60.
- [18] K. Nishimoto and Susana S. de Romero ; Numerical Examinations for Mixed and Double Finite Sums obtained by Means of N- Fractional Calculus, J. Frac.Calc.Vol. 26, Nov. (2004), 91 - 99.
- [19] K. Nishimoto ; Some Doubly Infinite, Finite and Mixed Sums in N- Fractional Calculus, J. Frac.Calc.Vol. 27, May (2005), 11 - 20.
- [20] Shy- Der Lin, H. M. Srivastava and Shih- Tong Tu ; Basic (or q -) Extensions of Some Multiple Infinite Sums Derived by Means of Fractional Calculus, Frac.Calc.Vol. 27, May (2005), 43- 57
- [21] K. Nishimoto ; Multiply Infinite Sums in N- Fractional Calculus and that in R - L One (Continue) Frac.Calc.Vol. 27, May (2005), 59 - 63.
- [22] K.S. Miller and B. Ross ; An Introduction to The Fractional Calculus and Fractional Differential Equations, John Wiley & Sons, (1993).
- [23] Igor Podlubny ; Fractional Differential Equations (1999), Academic Press.
- [24] R. Hilfer (Ed.) ; Applications of Fractional Calculus in Physics, (2000), World Scientific, Singapor, New Jersey, London, Hong Kong.
- [25] A.P. Prudnikov, Yu. A. Bryckov and O.I. Marichev ; Integrals and Series, Vol. I, Gordon and Breach, New York, (1986).
- [26] S. Moriguchi, K. Udagawa and S. Hitotsumatsu ; Mathematical Formulae, Vol.2, Iwanami Zensho, (1957), Iwanami, Japan.

Katsuyuki Nishimoto
 Institute of Applied Mathematics
 Descartes Press Co.
 2 - 13 - 10 Kaguike, Koriyama
 963 - 8833 Japan

Susana S. de Romero
 and Ana I. Prieto
 C.I.M.A., Universidad del Zulia
 Apartado 10482, Maracaibo
 Venezuela